GENERAL INDEX TO VOLUME XLI

New scientific names of plants and the final members of new combinations are printed in **bold face** type; synonyms and page numbers having reference to figures and plates, in *italics;* and all other matter in ordinary type.

A

Acerates, 47, 169; *angustifolia,* 181; *asperula,* 193; *atropurpurea,* 168; *auriculata,* 186; *circinalis,* 196; *connivens,* 160; *cordifolia,* 168; *decumbens,* 195; *delticola,* 172; *feayi,* 163; *floridana,* 172; *gomphocarpoides,* 189; *hirtella,* 170; *humilis,* 111; *incarnata,* 172; *lanceolata,* 173; *lanuginosa,* 175; *latifolia,* 180; *linearis,* 173; *longifolia,* 171; *monocephala,* 175; *nuttalliana,* 175; *obovata,* 141; *paniculata,* 191; *Pringlei,* 111; *rusbyi,* 183; *tomentosa,* 177; *vinosa,* 169; *viridiflora,* 173, var. *lanceolata,* 173, var. *linearis,* 173, var. *ivesii,* 173

Acerotis, 47

Adenostoma, Introgression in, 339

Adenostomma fasciculatum, 339, var. obtusifolium, 349

Agricultural practices in Honduras, 213, *218*

Alborotos, 220, *220*

Amaranth seeds, confections from, 220

Anantherix, 47, 160; *angustifolia,* 195; *connivens,* 160; *decumbens,* 195; *grandiflora,* 191; *nuttalliana,* 195; *ovata,* 191; *paniculata,* 191 *pumilis,* 199; *torreyana,* 191; *viridis,* 191

Anderson, Burton R.: A study of American petrified calamites, 395; Edgar Anderson and: Introgression of Salvia apiana and Salvia mellifera, 329

Anderson, Edgar: An analysis of introgression in a population of stemless white violets, 263; A field survey of chromosome numbers in the species of Tradescantia closely allied to Tradescantia virginiana, 305; Introgression in Adenostoma, 339; and Burton R.: Introgression of Salvia apiana and Salvia mellifera, 329; and William L. Brown: Ladyfinger and Tom Thumb, two old varieties of popcorn, 301; and Louis O. Williams: Maize and sorghum as a mixed crop in Honduras, 213

Andropogon sorghum, 292

Anthanotis, 47; *pedicellata,* 199; *procumbens,* 171, 191; *viridis,* 191

Arborescent lycopod fructifications, Some American, 351

Armenia, maize plants from, *286,* 287

Arthropitys, 398; bistriata, 395, 402, 413; communis, 395, 397, 401, 413, var. septata, 401, 406, 413; gigas, 396, 401, 413; Hirmeri, 401, 413; **illinoensis,** 398, 413, *415, 416;* kansana, 400, 402, 404, 413; sp., 401, *415; sp. B,* 399; **versifoveata,** 399, 413, *414, 415*

Arthroxylon, 398; *Williamsonii,* 402, 413, *415*

Asclepiadaceae, 1

Asclepias, L., The North American species of, 1; A correction in, 261

Asclepias, 1; economic uses of, 43; evolution in, 21; geographical distribution of, 27, *28,* 30, 34, 35; morphology of, 2, flower, 8, fruit, 18, inflorescence, 3, leaves, 3; pollination, 15, *44;* taxonomy, 47

Asclepias, 47; *aceratoides,* 124, 136; *acornutum,* 168; *acuminata,* 80; albicans, 147; *albiflora,* 51; *alticola,* 84; *amoena,* 51, 118, 156; amplexicaulis, 91; *amplexicaulis,* 108; *amsonioides,* 90; angustifolia, 62; *angustifolia,* 62, *128; apocinum,* 105; *apocynifolia,* 83; arenaria, 138; *arenicola,* 124; **asperula,** 193, ssp. asperula, 195, ssp. **capricornu,** 195; **atroviolacea,** 198; *aurantiaca,* 59; auriculata, 153; *auriculata,* 184; *bartlettiana,* 158; *bicolor,* 59; *bicknellii,* 87; *bidentata,* 84; brachystephana, 164; *brasiliensis,* 151; *brevicornu,* 156; californica, 177, ssp. californica, 178, ssp. **greenei,** 178; *capitellata,* 105; *capricornu,* 195, ssp. *occidentalis,* 195; *chapalensis,* 133; cinerea, 62; circinalis, 196; *citrifolia,* 116; *compressa,* 118; connivens, 160; *constricta,* 111; contrayerba, 86; **conzattii,** 129; *cordata,* 80; cordifolia, 168; *cornuti,* 105; coulteri, 89; **crocea,** 95; cryptoceras, 178, ssp. cryptoceras, 180, ssp. **davisii,** 180; *cubensis,* 59; curassavica, *5, 45, 46,* 59, var. *concolor,* 59; curtissii, 124; *curvipes,* 120; cutleri, 166; *dasypus,* 118; *davisii,* 180; *debilis,* 55; *decumbens,* 76, 195; *demissa,* 143; *douglasii,* 124; *eastwoodiana,* 165; elata, 186; *elliptica,* 105; emoryi, 158; engelmanniana, 184, var. *rusbyi,* 183; eriocarpa, 144, var. *microcarpa,* 144; erosa, 143; euphorbiaefolia, 126; exaltata, 87; *exilis,* 94; fascicularis, 64; *fasciculata,* 64; feayi, 163; *filiformis,*

113; *floridana*, 76; fournieri, 189; *fragrans*, 105, 153; *fremontii*, 144; fruticosa, 151; *galioides*, 67; gentryi, 73; *glaberrima*, 97; *gladewitzii*, 91; glaucescens, 97, var. *elata*, 186; *gonialis*, 118; grandiflora, 96; *grandifolia*, 105; *grayi*, 189; *guatemalensis*, 84; hallii, 120; hirtella, 170; humistrata, 108; *hybridus*, 116; hypoleuca, 110; incarnata, 50, ssp. incarnata, 51, ssp. pulchra, 53, var. *longifolia*, 51, var. *neoscotica*, 53, var. *pulchra* f. *candida*, 53, β. *pulchra*, 53, f. *albiflora*, 51; *incarnata*, 171; *insignis*, 96; *intermedia*, 105; involucrata, 130, var. *tomentosa*, 131; *ivesii*, 173; *jaliscana*, 62, 86; *jamesii*, 139; *kansana*, 105; *kotolo*, 144; labriformis, 142; lanceolata, 81; *lanceolata*, 173, var. *paupercula*, 81, f. *flaviflora*, 81; *lancifolia*, 55; lanuginosa, 121; *lanuginosa*, 261; *lasiotis*, 53, 118; latifolia, 139; *laurifolia*, 80; laxiflora, 134; lemmoni, 133; leptopus, 72; *leucophylla*, 143; linaria, 113; *lindheimeri*, 156; linearis, 58; *linifolia*, 62; *lonchophylla*, 120; *longicornu*, 156; longifolia, 171; *longifolia*, 62; *longipedunculata*, 93; *longipetala*, 191; *lutea*, 76; *macrophylla*, 64, var. *comosa*, 64; macrosperma, 131; macrotis, 132; *macroura*, 186; *margaritacea*, 59; *maritima*, 53; masonii, 145; *mazatlanensis*, 153; mcvaughii, 197; *meadii*, 109; *megalotis*, 136; *melantha*, 134; mexicana, 65, 67; michauxii, 128; mirifica, 188; *neglecta*, 83; nivea, 60, var. *curassavica*, 59, var. *intermedia*, 60, β. *flava*, 60, α. *normalis*, 60; nummularia, 149; *nutans*, 173; nuttalliana, 175; *nuttalliana*, 241; nyctaginifolia, 155; obovata, 141; *obtusata*, 143; *obtusifolia*, 91, β. *latifolia*, 39; oenotheroides, 156; otarioides, 261; *otarioides*, 121; ovalifolia, 104; ovata, 83; *palmeri*, 90; *parviflora*, 55, 68, var. *latifolia*, 55; *parvula*, 54; *paupercula*, 81; pedicellata, 199; pellucida, 103; perennis, 55, var. *parvula*, 54; *periplocaefolia*, 80; *phenax*, 111; *phytolaccoides*, 87; *pinifolia*, 113; *plumerifolia*, 153; *polyphylla*, 97; *polystachia*, 87; pratensis, 112; pringlei, 111; *procumbens*, 191; pseudorubricaulis, 59; puberula, 132; *pubescens*, 105; *pubigera*, 105; *pulchella*, 55; *pulchra*, 53, f. *albiflora*, 53; pumila, 71; purpurascens, 118; *purpusii*, 153; *quadrifolia*, 199, var. *oppositifolia*, 100; quinquedentata, 90, var. *neomexicana*, 90; *rafaelensis*, 94; *rolfsii*, 77; rosea, 150; *rothrockii*, 143; *rotundifolia*, 91; rubra, 80, var. *laurifolia*, 80; *rubricaulis*, 62; rusbyi, 183; ruthiae, 165; *salicifolia*, 62; scaposa, 94; *schaffneri*, 90; *scheryi*, 121; *senecionifolia*, 150; *serica*, 105; *serpentaria*, 81; *setosa*, 86; similis, 84; *simpsoni*, 163; solanoana, 180; speciosa, 124; sperryi, 190; *standleyi*, 160; stenophylla, 181; subaphylla, 146; subulata, 154; subverticillata, 65; *suffrutex*, 72; sullivantii, 115; syriaca, 44, 105, var. *kansana*, 105, β. *exaltata*, 87, f. *inermis*, 105, f. *leucantha*, 105; *tapalpana*, 62; texana, 54; *tithymaloides*, 89; tomentosa, 136 tuberosa, 7, 74, ssp. interior, 77, ssp. rolfsii, 77, ssp. tuberosa, 76, ssp. terminalis, 79, var. *flexuosa*, 77, var. *interior*, 77, var. *Rolfsii*, 77, β. *decumbens*, 76, f. *bicolor*, 77, f. *flavescens*, 77, f. *lutea*, 77; uncialis, 167; *vanillea*, 100; variegata, 116, α. *major*, 116, β. *minor*, 104; *verecunda*, 51; verticillata, 68, var. *linearis*, 58, var. *pumila*, 71, var. *subverticillata*, 65, var. α. *galioides*, 67, var. β. *mexicana*, 67; vestita, 148, ssp. parishii, 149, ssp. vestita, 149, var. *parishii*, 149; vinosa, 169; *virgata*, 62; viridiflora, 173, var. *linearis*, 173, β. *lanceolata*, 173, γ. *obovata*, 141; viridis, 191; viridula, 102; virletii, 93; woodsoniana, 57; *wrightii*, 155; zanthodacryon, 190

Asclepiodella, 47, 161; *feayi*, 163

Asclepiodora, 47, 185; *asperula*, 193; *circinalis*, 196; *decumbens*, 195; *feayi*, 163; *gomphocarpoides*, 189; *insignis*, 96; *virdis*, 191; *zanthodacryon*, 190

"Asia, Some peculiarities of maize in," by Kuleshov, translation of, 271

Asterophyllites charaeformis, 403

Astromyelon, 404; augustodunense, 406; cauloides, 405, 413, *414*, *416*; pluriradiatum, 406, 413, *415*, *417*; reticulatum, 406; Williamsonis, 399, 413

Atolillos, 221

Autoploidy, 305

Azerbaijan, maize plants from, 287, *288*

B

Biventraria, 47; *variegata*, 116

Blepharodon gomphocarpoides, 189

Bothrodendron, 375

Brown, William L., Edgar Anderson, and: Ladyfinger and Tom Thumb, two old varieties of popcorn, 301

C

Caesalpiniaceae, 223
Calamites, A study of American petrified, 395
Calamites, *multifolia*, 408; **rectangularis**, 403, 413, *416*
Calamodendron, 398
Calamospora, 409
Cantheliophorus, 380
Cardiocarpon, 381
Calabash tree fruits as seed pouches, 214, *219*
Carboniferous fossil floras, fructifications of, 351, 395
Chamise, 339
China, western, types of maize plants from, 272, *276*
Chromosome numbers in species of Tradescantia closely allied to Tradescantia virginiana, 305, 322
Coal balls, American, 351, 395
Collins' work on American maize, 272, 298
Condensation in inflorescence of Adenostoma, 346
Cone genera, fossil, 351, 395, 407
Corn, 274; see also Maize and Zea
Crescentia alata, 213; cujete, 213
Cuba, 226
Cystosporites, 379, 380, 382; giganteus, 368

D

Dicymbe, 254; **psilophylla**, 254
Diploid Tradescantias, distribution of, *321*, 322–324
Dwyer, John D.: The tropical American genus Tachigalia Aubl., 223

E

Ecological dominants in population of Salvia apiana and S. mellifera, 337
Economic uses of Asclepias, 43
Equisetum, 402, 410
Evolution: in Adenostoma, 350; in Asclepias, 21; in Lepidostrobus, 379

F

Felix, Charles J.: Some American arborescent lycopod fructifications, 351
Field survey of chromosome numbers in the species of Tradescantia closely allied to Tradescantia virginiana, 305
Fossil fructifications, 351, 395

G

Geographical distribution of North American species of Asclepias, 27, *28*, 30, *34–35*
Geography of races of maize, 271
Gomphocarpus, 47; ***angustifolius***, 151; *arachnoideus*, 151; *auriculatus*, 185; *brasiliensis*, 151; *cordifolius*, 168; *fruticosus*, 151, f. *brasiliensis*, 151; *hypoleucus*, 110; *longifolius*, 172; *palmeri*, 189; *purpurascens*, 180; *tomentosus*, 177, var. *xanti*, 178; *torreyi*, 177, var. *xanti*, 178; *viridiflorus*, 173; *viridis*, 191
Guatemala, Notes on sorghum in, 220

H

Honduras; Maize and sorghum as a mixed crop in, 213; planting operations, *218*; maize in field, *219*; variation of maize in, *216*, *219*, pls. *7*, *8*
Hybridization: in Asclepias, 23; in Salvia, 329; in Tradescantia, 307

I

Illiniocarpon, 379, 382
Inflorescence variation in Adenostoma, 339, *340–343*, *354*
Insects: ants inhabiting species of Tachigalia, 224; pollination by, in Asclepias, 14, 16, *44*, in Salvia, 329
Introgression: in Adenostoma, 339; in a population of stemless white violets, 263; of Salvia apiana and Salvia mellifera, 329; in species of Tradescantia closely allied to T. virginiana, 305
Isoetes, 364

J

Jicaras, *219*

K

Kharkov, maize plants grown at, 272, *276*, *282*, *283*, *290*
Kempton's maize investigations, 271
Kidd, Harold J., and Howard C. Reynolds: Translation of Kuleshov's "Some peculiarities of maize in Asia", 271
Kuleshov's "Some peculiarities of maize in Asia", translation of, 271

L

Ladyfinger and Tom Thumb, two old varieties of popcorn, 301, *304*
Lagenicula rugosa, 359; saccata, 368, 380
Leaves, Calamitean, 402
Lebrunniodendron, 224

Lepidocarpon, 351, 374, 379, 381; **crenatum**, 385, *394*; ioense, 382; Lomaxi, 379; magnificum, 382, *383*, *393*
Lepidodendron, 356, 363, 371, 375, 376; acutum, 375; Brownii, 376; obovatum, 376; simile, 375; Veltheimianum, 375
Lepidophloios, 374, 379
Lepidostrobus, 351; allantonensis, 375; *Arberi*, 374; *aristatus*, 366; arrectus, 366; Bartletti, 367; Binneyanus, 358, 374; Bohdanowiczii, 379; braidwoodensis, 368, 376, 379; Brownii, 353, 360, 372, 379; coulteri, 353, 368, 372, 374, 376, 382; **diversus**, 353, *354*, *361*, *362*, *388*, *389*, *390*; dubius, 359, 370, 375, 378, f. hermaphroditus, 378, f. megalophorus, 378; *Fischeri*, 371; *foliaceous*, 377; Gallowayi, 367, 369, 379; gracilis, 374; imbricatus, 357, 371; Jacksoni, 376; kentuckiensis, 371; major, 378, 380; Masleni, 356, 376; monospora, 380; noei, 357, 360, 373, 379, *390*; oldhamius, 374, 376, 382, f. pilosus, 374; olryi, 359, 375; **pulvinatus**, 362, *365*, *391*, *392*; russelianus, 353, 359, 375, 378; scleroticum, 361; *Scottii*, 374; sp., 368; Veltheimianus, 374; zea, 376, 379
Ligule in Lepidostrobus, 374
Liguleless maize, 293, from Primorskiĭ Province, 297
Lycopod fructifications, Some American arborescent, 351
Lycopodiaceous strobilus, A, 369
Lycospora, 360, 366, 370, 382; micropapillatus, 366; parva, 366; punctata, 382; rugosa, 361

M

Maicillo, 221
Maize: American, grown in Asia, 272, 276; Armenian, *286*, 287; Asiatic, 272; brachytic, 272, *275*; Chinese, 272, *276*; flint, 274; Kharkov, 272; in Honduras, 213; liguleless, 293, *297*; Longfellow, 279, 281; Manchurian, 272, *276*; Mexican, 215; Minnesota, 279, 281; Persian type, 279, *280*, *283*, *284*; Primorskii, 272; Turkestan, 284, *286*; variation in Honduran, *216*, *219*, *pls. 7*, *8*; waxy, 272, *273*, *278*; Zapaluta chica, 215
"Maize: Some Peculiarities of, in Asia", by Kuleshov, translation of, 271
Maize and sorghum: as a mixed crop in Honduras, 213, *218*, *219*; convergant appearance of, from same regions, 291
Man as an ecological dominant in populations of Adenostoma, 350
Manchuria, maize plants from, 272, *276*
Mass collections of stemless white violets, 263
Mazocarpon, 351, 364, 376, 380, 412; oedipternum, 370, 380; pettycurense, 381; shorense, 375, 381
Myriophylloides, 405
Myrmecophily in Tachigalia, 224

N

North American maize grown in Asia, 272, 276
North American species of Asclepias L., 1

O

Oligoron, 47; *longifolium*, 172, var. *birsutum*, 170; *tenuifolium*, 128
Otanema, 47; *lanceolata*, 173; *latifolia*, 139, 173; *ovata*, 173
Otaria, 47; *auriculata*, 153; *fragrans*, 153; *laxiflora*, 134
Oxypteryx, 47; *arenicola*, 124; *curtissii*, 124

P

Paleobotany: Some American arborescent lycopod fructifications, 351; A study of American petrified Calamites, 395
Paleostachya, 407; arborescens, 407; multifolia, 408, *411*, 413, *417*, *418*; Schimperiana, 407; sp., 411; vera, 410
Panicum italicum, compared with sorghum and maize in habit, *294*, 295
Persian type of mazie, 279, *280*, *283*; vegetative diversity of, *284*
Petrified calamites, American, A study of, 395
Phenological observations on maize at Kharkov, 277, 282
Plani-sporites, 369, 381
Platymiscium, 224
Podostemma, 47; *australe*, 156; *emoryi*, 158; *helleri*, 156; *leoninum*, 158; *Lindheimeri*, 156; *longicornu*, 156; *nyctaginifolium*, 155
Podostigma, 47, 152, 196; *pedicellata*, 199; *pubescens*, 199; *viridis*, 191
Pollination, insect, in Asclepias, 14, *44*; in Salvia, 329
Polyotus, 47, 181; *angustifolius*, 181; *heterophyllus*, 173; *lanuginosus*, 175; *longifolius*, 172; *obovatus*, 141
Polyploidy, 305
Popcorn, Ladyfinger and Tom Thumb, two old varieties of, 301, *304*
Popcorns of Guatemala, 200
Population studies: in Adenostoma, 347; of Salvia apiana and S. mellifera, 332

Porostrobus, 375
Primorskiĭ Province, liguleless maize from, 297; waxy maize grown in, 277, *278*

R

Reynolds, Howard C., Harold J. Kidd and: Translation of Kuleshov's "Some peculiarities of maize in Asia", 271
Rio Yeguare, Honduras maize and sorghum in valley of, 213; samples of maize planted in, *219*, *pls. 7, 8*
Roots, petrified calamitean, 404

S

Sages, 329
Salvia apiana var. compacta, 333
Salvia: hybridization in, 332; morphological analysis of species differences in, 329; variation pattern produced by introgression in, *330*, *331*, 333, 334
San Gabriel Mountains, introgression between Salvia apiana and Salvia montana in, 332
Sauer, Jonathan: Notes on Sorghum in Guatemala, 220
Schizonotus, 47, 176; *purpurascens*, 181
Sclerolobium, 223; myrmecophilum, 244; odoratissimum, 224; physophorum, 224
Selaginella, 378; Vögelii, 378
Sigillaria, 360, 380
Sigillarian fructifications, 330
Sigillariostrobus, 351, 376, 378, 380; ciliatus, 380; Czarnockii, 380; rhombibracteatus, 369, 380; sphenophylloides, 380
Solanoa, 47, 177; *purpurascens*, 180
Solanoana, 177; *purpurascens*, 180
Sorghum: maize and, as a mixed crop in Honduras, 213; plants from Asia compared, 291, *295*; notes on, in Guatemala, 220; planting, *218*; in field, *219*; stems of, with leaf sheaths removed, *296*
Sorghum caffrorum var. glabrescens in Honduras, 215; vulgare, 221
Stems, petrified, 395, 401
Stratigraphic, correlations of coal ball impressions, 412
Strobilus, A lycopodiaceous, 369
Stylandra, 47; *pumila*, 199

T

Tachia, 226
Tachigalia Aubl., the tropical American genus (Caesalpiniaceae), 223; myrmecophily in, 224; taxonomy, 226
Tachigalia, 226; *alba*, 238, 240; agyrophylla, 247, *260*; *angustifolia*, 238, 239; **bracteolata**, 233; *carinata*, 238; catingae, 229, *256*; cavipes, 234, *256*; *eriocalyx*, 238; formicarum, 245, *258*; grandiflora, 254; grandistipulata, 243, *258*; longiflora, 228, *256*; macrostachya, 251; multijuga, 230, *256*; myrmecophila, 244, *258*; paniculata, 238, *256*, var. δ. **alba**, 240, *258*, var. α. **angustifolia**, 239, *258*, var. *cavipes*, 234, var. γ. **comosa**, 240, *258*, var. β. **sulcata**, 240, *258*; plumbea, 242, *258*; *poeppigiana*, 246; polyphylla, 246, *260*; *psilophylla*, 254; ptychophysca, 252, *260*; pubiflora, 236, *256*; **pulchra**, 249, *260*; rigida, 232, var. argentea, 233; rusbyi, 248, *260*; *sericea*, 238; *sulcata*, 238, 240; tessmannii, 250, *260*; *trigona*, 238; ulei, 241, *258*; **venusta**, 235, 256; versicolor, 253, *260*
Tassia, 226
Tetraploid and diploid Tradescantias, maps showing distribution of, *306–321*
Tom Thumb, Ladyfinger and, Two old varieties of popcorn, 301
Tradescantia, Chromosome numbers in species of, closely allied to Tradescantia virginiana, 305
Tradescantia, 305; bracteata, 315, *316*, 325; edwardsiana, 308, *308*, 322; ernestiana, *309*, 310, 323; gigantea, 312, *313*, 324; hirsuticaulis, 316, *317*, 326; hirsutiflora, 311, *311*, 323; humilis, 318, *319*, 326; longipes, 316, *317*, 326; occidentalis, 320, *321*, 326; ohiensis, 313, *314*, 324; ozarkana, 308, *309*, 322; paludosa, 314, *315*, 325; reverchonii, 318, *318*, 326; roseolens, 320, *320*, 326; subacaulis, 318, *319*, 326; subaspera var. typica, 306, *306*, 322, var. montana, 307, *307*, 322; tharpii, 312, *312*, 323; virginiana, 310, *310*, 323
Tradescantia virginiana group of plants, chromosome counts of, 305, 323
Triletes, 358, 368, 373; auritus, 376; diabolicus, 377; glabratus, 380; horridus, 359, 378; mamillarius, 380; rugosus, 358, 375; translucens, 358
Tropical American genus Tachigalia Aubl. (Caesalpiniaceae), 223
Turkestan collections of maize, 285, *286*
Twigs, Calamitean, 402

V

Valentina, 226
Variation in Adenostoma, 339; in Honduran maize, 215, *216*, *219*, pls. *7*, *8*; in

petrified stems, 395; in Salvia apiana and S. mellifera, *331*, 334, *335*; in Viola pallens, 263, *265*, *267*; in maize, 272

Vavilov's work on diversity of crop plants, 298

Viola blanda, 268; incognita, 268; pallens, introgression in, 263

Violets, stemless white, An analysis of introgression in a population of, 263

W

Wasp visiting Asclepias syriaca, *44*

Williams, Louis O., Edgar Anderson and: Maize and sorghum as a mixed crop in Honduras, 213

Woodson, Robert E., Jr.: A correction in Asclepias, 261; The North American species of Asclepias L., 1

Z

Zea mays, 301, var. certina, 278